

UNCLASSIFIED

AD NUMBER
AD813720
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Critical Technology; APR 1967. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab ltr dtd 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

TECHNICAL MANUSCRIPT 376

INHIBITION OF FLOWERING IN
XANTHIUM PENSYLVANICUM WALLN.
BY ETHYLENE

Frederick B. Abeles

STATEMENT OF UNCLASSIFIED

This document is subject to special export controls and each transmission to the government of foreign nationals may be made only with prior approval of

DA
Ft. Detrick
Frederick, Md.


APRIL 1967

MAY 15 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

AD 813720

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 376

INHIBITION OF FLOWERING IN XANTHIUM PENSYLVANICUM WALLN.
BY ETHYLENE

Frederick B. Abeles

Crops Division
BIOLOGICAL SCIENCES LABORATORY

Project 1L013001A91A

April 1967

INHIBITION OF FLOWERING IN XANTHIUM PENSYLVANICUM WALLN.
BY ETHYLENE

ABSTRACT

Ethylene inhibited the formation of flowers in cocklebur (Xanthium pensylvanicum Walln.) plants that were exposed to an inductive dark period. Indoleacetic acid had been shown earlier to have this same effect. Because it stimulates ethylene evolution from cocklebur leaves, the results suggest that the effect of auxin may be an ethylene effect.

Bonner and Thurlow¹ reported that indoleacetic acid (IAA) inhibited the formation of flowers in cockleburs (Xanthium pensylvanicum Walln.) when it was applied as a spray during the long inductive night. In the present report, evidence is presented that ethylene also inhibits the flowering of cocklebur plants that have been exposed to a long inductive photoperiod and that the IAA effect is probably due to auxin-stimulated ethylene production.

The two large apical beaks were cut from cocklebur fruits (Chicago strain*) and the fruits were washed in running water for 24 hours and planted in 4-inch pots of soil; seedlings were grown for 3 weeks at 25 C on a 16-hour light cycle (1,500 ft-c) supplied by daylight fluorescent bulbs. The plants were induced to flower by exposure to 16 hours of darkness in 10-liter desiccators containing 0, 1, 10, or 100 ppm ethylene. After the induction period, the plants were returned to the growth chamber; the sizes of the inflorescences were measured 8 weeks later. Ethylene evolution from the plants was measured by gas chromatography.²

The inflorescences of the two controls receiving no ethylene were 5 and 8 mm in diameter and in anthesis. After 16 hours in the desiccator, the level of ethylene in the gas phase had risen to 0.025 ppm. One of the plants treated with 1 ppm ethylene remained vegetative and the other had an inflorescence 2 mm in diameter. The two plants treated with 10 and 100 ppm ethylene remained vegetative. Except for an initial temporary leaf epinasty, the ethylene treatment had no effect on the subsequent growth and vigor of the plants when compared with controls maintained on long days. This experiment was repeated on three other occasions and

* Supplied by F.B. Salisbury, Utah State University, Logan, Utah.

similar results were obtained. Ethylene production from untreated cocklebur plants during the 16-hour dark period was 7 nl per hour per plant. Spraying prior to the dark period with 10^{-4} and 10^{-3} M IAA until runoff increased the rate of gas production to 14 and 28 nl per hour, respectively.

Some recent examples of auxin effects shown to be ethylene effects include the stimulation of abscission by auxin,² the auxin-induced inhibition of growth of pea stem sections,³ and the induction of flowering of pineapples.⁴ Palmer⁵ recently reported that high levels of IAA would inhibit the induction of invertase in aged beetroot (Beta vulgaris L.) tissue slices. Using the same species Scott⁶ reported that 10 ppm ethylene would also inhibit invertase synthesis. Another example of auxin-ethylene effects may be the development of sex in monoecious species. In a recent review, Nitsch⁷ pointed out that treatments by auxin and unsaturated gases such as ethylene, CO, and acetylene favor the formation of female flowers on cucumber (Cucumis sativus L.) plants. The results presented here suggest that the inhibition of flowering in the cocklebur by auxin may also be an ethylene effect.

The mechanism by which ethylene prevents the induction of floral primordia is unknown. However, it is known that RNA synthesis^{8,9} and presumably protein synthesis^{8,10} are an essential part of the induction process. It is known that ethylene can enhance RNA and protein syntheses in ethylene-sensitive tissue,¹¹ suggesting that ethylene may interfere in some way with the orderly production of RNA required for the flowering process.

LITERATURE CITED

1. Bonner, J.; Thurlow, J. 1949. Inhibition of photoperiodic induction in Xanthium by applied auxin. Bot. Gaz. 110:613-624.
2. Abeles, F.B.; Rubinstein, B. 1964. Regulation of ethylene evolution and leaf abscission by auxin. Plant Physiol. 39:963-969.
3. Burg, S.P.; Burg, E.A. 1966. The interaction between auxin and ethylene and its role in plant growth. Proc. Nat. Acad. Sci. 55: 262-269.
4. Burg, S.P.; Burg, E.A. 1966. Auxin-induced ethylene formation: Its relation to flowering in the pineapple. Science 152:1269.
5. Palmer, J.M. 1966. The influence of growth regulating substances on the development of enhanced metabolic rates on thin slices of beetroot storage tissue. Plant Physiol. 41:1173-1178.
6. Scott, P.C. 1966. Abscission as a cellular senescence. Ph.D. Thesis. Purdue University, Lafayette, Indiana.
7. Nitsch, J.P. 1965. Physiology of flower and fruit development. In A. Lang (ed.) Encyclopedia of plant physiology. Springer-Verlag Berlin. 15, part 1:1537-1647.
8. Collins, W.T.; Salisbury, F.B.; Ross, C.W. 1963. Growth regulators and flowering: III. Antimetabolites. Plants 60:131-144.
9. Salisbury, F.B.; Bonner, J. 1960. Inhibition of photoperiodic induction by 5-fluorouracil. Plant Physiol. 35:173-177.
10. Miller, J.; Ross, C.W. 1966. Inhibition of leaf processes by p-fluorophenylalanine during induction of flowering in the cocklebur. Plant Physiol. 41:1185-1192.
11. Abeles, F.B.; Holm, R.E. 1966. Stimulation of RNA synthesis, protein synthesis, and abscission by ethylene. Plant Physiol. 10:1337-1342.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author)		2a REPORT SECURITY CLASSIFICATION
DEPARTMENT OF THE ARMY Fort Detrick, Frederick, Maryland 21701		Unclassified
		2b GROUP
3 REPORT TITLE		
INHIBITION OF FLOWERING IN <u>XANTHIUM PENSYLVANICUM</u> WALLN. BY ETHYLENE		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5 AUTHOR(S) (Last name, first name, initial)		
Abeles, Frederick B.		
6 REPORT DATE	7a TOTAL NO. OF PAGES	7b NO. OF REFS
April 1967	8	11
8a CONTRACT OR GRANT NO.	9a ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO. 1L013001A91A	Technical Manuscript 376	
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		
10 AVAILABILITY/LIMITATION NOTICES		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11 SUPPLEMENTARY NOTES		12 SPONSORING MILITARY ACTIVITY
		Department of the Army Fort Detrick, Frederick, Maryland 21701
13 ABSTRACT		
Ethylene inhibited the formation of flowers in cocklebur (<u>Xanthium pensylvanicum</u> Walln.) plants that were exposed to an inductive dark period. Indoleacetic acid had been shown earlier to have this same effect. Because it stimulates ethylene evolution from cocklebur leaves, the results suggest that the effect of auxin may be an ethylene effect.		
14. Key Words		
<u>Xanthium pensylvanicum</u> Ethylene Auxins Inhibiting Flowering		

DD FORM 1 JAN 64 1473

Unclassified
Security Classification